

SMALL 3-MANIFOLDS OF LARGE GENUS

IAN AGOL

ABSTRACT. We prove the existence of pure braids with arbitrarily many strands which are small, *i.e.* they contain no closed incompressible surface in the complement which is not boundary parallel. This implies the existence of irreducible non-Haken 3-manifolds of arbitrarily high Heegaard genus.

1. INTRODUCTION

Haken introduced the notion of irreducible 3-manifolds containing an incompressible surface, called sufficiently large, or Haken [6]. These were thrust into prominence by theorems of Waldhausen, who (among other things) showed that the homeomorphism and word problems are solvable for these manifolds [21], and by Thurston who showed that they satisfy the geometrization conjecture [20]. It is therefore interesting to understand how prevalent non-Haken 3-manifolds are. Jaco and Shalen showed that a Seifert fibred space M is non-Haken if and only if it is atoroidal with $H_1(M)$ finite [11]. Thurston showed that all but finitely many Dehn fillings on the figure eight knot complement are non-Haken [19]. Hatcher and Thurston extended this result to all 2-bridge knot exteriors [9]. These examples all have Heegaard genus 2. By a result of Hatcher, if a link is small, *i.e.* is irreducible and contains no closed incompressible surface other than boundary tori, then Dehn filling on any boundary component yields a small manifold for all but finitely many fillings [10]. Floyd and Hatcher [5] and independently Culler, Jaco, and Rubinstein [3] showed that punctured torus bundles and 4-punctured sphere bundles with irreducible monodromy are small. Oertel proved that Montesinos knots covered by small Seifert fibred spaces are small [17]. It is believed that the Seifert-Weber dodecahedral space is small. Several other examples of small manifolds are known, *e.g.* the Borromean rings (Lozano), some chain links with ≤ 5 components (Oertel), and some other examples of Dunfield [4], Hass and Menasco [7], and Lopez [12, 13].

A result of Moriah and Rubinstein shows that the Heegaard genus of infinitely many Dehn fillings on a link complement is the same as that of the link complement [16]. Since most punctured torus bundles have Heegaard genus 3 [2], there are infinitely many closed small 3-manifolds of genus 3. This appears to be the largest known genus of a small 3-manifold in the literature. Alan Reid asked whether there are small links of arbitrarily many components, observing that this would imply the existence of irreducible non-Haken 3-manifolds of arbitrarily large Heegaard genus, as we show in theorem 3.2. We prove in theorem 3.1 the existence of small links of arbitrarily many components, answering Reid's question.

Acknowledgments. We thank Alan Reid, Hyam Rubinstein, Saul Schleimer, and Bill Thurston for contributing ideas and suggestions which led to the results in this paper. The ideas in the paper were partly inspired by a talk of Dan Margalit at UIC on the pants complex [14], and by the work of Jeff Brock [1].

Date: February 1, 2008.


1991 Mathematics Subject Classification. Primary 57M50; Secondary 57M25.

Key words and phrases. Heegaard genus, 3-manifold, Haken.

Partially supported by ARC grant 420998.

2. PANTS DECOMPOSITIONS

The pants complex of a surface was introduced by Hatcher, Lochak, and Schneps [8], and we will follow their conventions and terminology. We will just be interested in the pants graph. Let Σ be a connected compact orientable surface with $\chi(\Sigma) < 0$. We say Σ has type (g, n) if it has genus g and n boundary components. By a **pants decomposition** of Σ , we mean a finite collection P of disjoint smoothly embedded circles cutting Σ into pieces which are surfaces of type $(0, 3)$. The number of curves in a pants decomposition is $3g - 3 + n$, and the number of pants is $-\chi(\Sigma)$. Let P be a pants decomposition, and suppose that one of the circles β of P is such that deleting β from P produces a complementary component of type $(1, 1)$. This is equivalent to saying there is a circle γ in Σ which intersects β in one point transversely and is disjoint from all the other circles in P . In this case, replacing β by γ in P produces a new pants decomposition P' (fig. 1). We call this replacement a **simple move**, or *S*-move. In a similar fashion, if P contains a circle β such that deleting β from P produces a complementary component of type $(0, 4)$, then we obtain a new pants decomposition P' by replacing β with a curve γ intersecting β transversely in two points and disjoint from the other curves of P (fig. 1). The transformation $P \rightarrow P'$ in this case is called an **associativity move**, or *A*-move.

FIGURE 1. An *A*-move and *S*-move

Definition 2.1. The **pants decomposition graph** $\mathcal{P}(\Sigma)^{(1)}$ is the graph having vertices the isotopy classes of pants decompositions of Σ , with an edge joining two vertices whenever the corresponding pants decompositions differ by a single *A*- or *S*-move. This is the one skeleton of the pants decomposition complex, which was proven to be connected in [8].

Main Construction: Suppose we have a homeomorphism $\psi : \Sigma \rightarrow \Sigma$, with mapping torus $T_\psi = (S \times I) / \{(x, 0) \sim (\psi(x), 1)\}$. Given a path $C = P_0 - P_1 - \dots - P_{m-1} - P_m \subset \mathcal{P}(\Sigma)^{(1)}$, $P_m = \psi(P_0)$, we get a sequence of circles $\beta_1, \dots, \beta_m$, where β_{i+1} is the circle in P_{i+1} replacing a circle in P_i (i is taken mod m). We assume that there is no circle β_j which is contained in all of the partitions P_i , $1 \leq i \leq m$. For each circle β_i , drill out a curve $B_i = \beta_i \times \{\frac{i}{m}\}$ in T_ψ . For each complementary region of P_i with boundary curves $\beta_i, \beta_j, \beta_k$ (where i, j, k might not be distinct), there is an embedded pants in T_ψ with boundary on the curves $B_i \cup B_j \cup B_k$ and interior disjoint from $\cup_i B_i$, and moreover, all of these pants may be chosen to have disjoint interiors (we pull apart the pants lying in Σ like an accordion). For an *A*- or *S*-move $P_i - P_{i+1}$, there is a complementary region bounded by two pants for an *S*-move, and four pants for an *A*-move, which we call an *A*- or *S*-region, respectively. Consider the link complement $M_C = T_\psi \setminus \mathcal{N}(\cup_i B_i)$. Then T_ψ is obtained from M_C by Dehn filling on the boundary components corresponding to the link $\cup_i B_i$. M_C is decomposed along pants into *A*- or *S*-regions.

Definition 2.2. A surface $(S, \partial S) \subset (M_C, \partial M_C)$ (where if $C = \emptyset$, then $M_C = T_\psi$) is **pairwise incompressible** if

1. each component of ∂S is either parallel to a component of $\partial \Sigma$ in ∂T_ψ or to a longitude of a component of $\partial \mathcal{N}(B_i)$,
2. if there is an annulus $A \subset M_C$ such that $\text{int} A \cap S = \emptyset$, one boundary component of A is in $\text{int} S$ and the other is in ∂M_C parallel to a component of $\partial \Sigma$ or a longitude of $\partial \mathcal{N}(B_i)$, then A is isotopic into S .

Lemma 2.3. $\text{int} M_C$ has a complete hyperbolic metric of finite volume. Any pairwise incompressible surface in M_C is a disjoint union of pants. Any pairwise incompressible surface in T_ψ isotopic into M_C is obtained by tubing pants in M_C .

Proof. The A - and S -regions may be regarded as pared manifolds [15], with pared locus consisting of annuli in the boundary which are regular neighborhoods of the curves from the pants decomposition. Each A -region may be decomposed into two ideal octahedra (fig. 2), and each S -region is decomposed into one ideal octahedron (fig. 3).




FIGURE 2. An A -region from 2 octahedra





FIGURE 3. An S -region from one octahedron

Thus, an A -region is obtained by doubling a checkerboard colored ideal octahedron along the dark faces. An S -region is obtained by folding pairs of dark faces of a checkerboard colored ideal octahedron together along common ideal vertices. This gives each A - and S -region a hyperbolic metric with totally geodesic boundary, and rank one cusps along the pared locus. We may glue these pieces together along geodesic 3-punctured spheres to get a hyperbolic structure on $\text{int} M_C$.

Suppose that we have an incompressible surface $(S, \partial S) \subset (M_C, \partial T_\psi)$, such that S is pairwise incompressible in T_ψ . We may assume that it intersects each pants in essential simple closed curves. Since closed curves in pants are boundary parallel, we may do surgery along annuli, to get a surface S' which is disjoint from the pants (fig. 4). Since M_C is acylindrical, we may assume that none of the surgeries produce annuli. Thus, each component of the surface S' lies in an A - or S -region. Since

FIGURE 4. Surgering S along an annulus A to obtain S'

an ideal octahedron is the same as a truncated tetrahedron with its edges drilled out (where dark faces of the octahedron correspond to faces of the tetrahedron), we may assume that the surface intersects each octahedron of the decomposition of A - and S -regions normally. A surface is normal if and only if it is incompressible in the complement of the 1-skeleton [18], so since there is no one skeleton, the surface must be normal. In the case of an A -region, the surface must be a double of a quadrilateral or a triangle. The doubles of triangles give the pants surfaces, and the doubles of the quadrilaterals have annular compressions to the pared locus, that is they are obtained by tubing together pairs of pants. A similar property holds for the S -regions. Thus, we may obtain every pairwise incompressible surface by tubing together pants. $\square$

Corollary 2.4. $\text{Vol}(T_\psi) \leq V_{\text{oct}}(2A + S)$, where Vol denotes the volume of the complete hyperbolic metric, V_{oct} denotes the volume of a regular ideal octahedron, A is the number of A -moves in the path C , and S denotes the number of S -moves.

Proof. Each A -region of M_C contributes $2V_{\text{oct}}$, and each S -region contributes V_{oct} . Then by a theorem of Thurston [19], $\text{Vol}(T_\psi) < \text{Vol}(M_C) = V_{\text{oct}}(2A + S)$. $\square$

Remark: This gives a tight upper bound for a theorem of Brock [1] relating volumes of mapping tori to pants distance of the monodromy.

3. SMALL LINKS

Theorem 3.1. *There exist pure braids in $S^2 \times S^1$ with arbitrarily many components and no closed incompressible surface in the complement.*

Theorem 3.2. *There are small 3-manifolds of arbitrarily large Heegaard genus.*

Proof. 3.2 The following argument was observed by Alan Reid. By theorem 3.1, there is a small link with n components, for n arbitrarily large. A link complement with n boundary components has Heegaard genus $g \geq n/2$, since there must be $\geq n/2$ boundary components lying to one side of a Heegaard surface, and therefore the genus of the Heegaard surface is $\geq n/2$. A result of Hatcher [10] shows that there are finitely many Lagrangian subspaces of the space of measured laminations on the boundary which consist of laminations which are the boundary of incompressible measured laminations. Moriah and Rubinstein [16] prove that for each boundary component, there are finitely many points and lines in Dehn filling space of the component, such that if a Dehn filling avoids these, then the Heegaard genus of the resulting manifold is the same as the Heegaard genus of the link complement. Combining these two theorems, we see that there are infinitely many Dehn fillings on the link which are small with genus $\geq n/2$, by doing Dehn fillings which avoid the slopes given by Moriah and Rubinstein's theorem and avoiding the boundary slopes of surfaces given by Hatcher's theorem. $\square$

Proof. 3.1 Let Σ_n be the n -punctured sphere, i.e. a surface of type $(0, n)$. We construct a closed path C in the pants graph $\mathcal{P}(\Sigma_n)^{(1)}$ and apply the Main Construction to obtain a sequence of loops in $\Sigma_n \times S^1$. Then we do very high Dehn twists about these loops to force any incompressible surface in the resulting braid to be isotopic into the complement of these loops (using Hatcher's theorem). By lemma 2.3, any pairwise incompressible surface is obtained by tubing together pants.

Then we analyze the ways of tubing pants together, and show that the resulting surfaces are always compressible in the braid complement obtained by any Dehn twists about these loops. Bill Thurston suggested considering manifolds which fiber over S^1 , Saul Schleimer suggested drilling out horizontal curves in a braid to try to find small braids by large Dehn twists, and Hyam Rubinstein suggested using links whose complements decompose along pants.

The first observation is that if there is a closed incompressible surface in T_ψ , then there is a pairwise incompressible surface, obtained by doing all possible compressions along compressing annuli.

We choose a closed path in the pants graph $\mathcal{P}(\Sigma_n)^{(1)}$, such that each pants is a twice punctured disk or a punctured annulus, when thought of as lying in $\Sigma_n \times S^1 / \subset S^2 \times S^1$. If this is true, then after tubing along the horizontal loops, the surface becomes a punctured sphere, torus, or klein bottle. If there is a 2-punctured disk, then the surface must be a punctured sphere. The only essential punctured sphere in a braid complement is the fiber surface. To see this, we take a cyclic cover of $S^2 \times S^1$, and lift the punctured S^2 to this cover, so that it lies between two fiber surfaces. But the only incompressible surfaces in the complement of the fiber are copies of the fiber. If copies of the fiber are tubed together, then the resulting surface is compressible. Thus, we may assume that the only pants that occur are punctured annuli.

Now we construct an explicit closed path in $\mathcal{P}(\Sigma_n)^{(1)}$. This path seems to be the simplest to construct with the property that each pants in the path is a punctured annulus or 2-punctured disk. Consider the punctures of Σ_n as lying in a great circle γ on S^2 . Cyclically label the gaps between punctures along γ $1, \dots, n$. For every pair of gaps (i, j) , there is a loop $\beta_{i,j}$ in S^2 which meets γ precisely twice at the gaps i and j . We form a path $C \subset \mathcal{P}(\Sigma_n)^{(1)}$. The initial pants decomposition is given by $P_0 = \{\beta_{1,3}, \beta_{1,4}, \dots, \beta_{1,n-1}\}$. We may take another pants decomposition $P_{n-3} = \{\beta_{2,4}, \beta_{2,5}, \dots, \beta_{2,n}\}$, which is obtained from P_0 by shifting each index by 1. There is a path of pants decompositions $C_0 = P_0 - P_1 - \dots - P_{n-3}$, where P_k is obtained by replacing the first k loops of P_0 with the first k loops of P_{n-3} , $k = 1, \dots, n-3$ (fig. 5). More generally, let $P_{j(n-3)} = \{\beta_{j+1,j+3}, \beta_{j+1,j+4}, \dots, \beta_{j+1,j+n-1}\}$, where indices are taken $(\text{mod } n)$. For $1 \leq k \leq n-3$, let $P_{j(n-3)+k}$ be obtained by replacing the first k loops of $P_{j(n-3)}$ with the first k loops of $P_{(j+1)(n-3)}$. Then we have a closed path $C = P_0 - P_1 - \dots - P_{n(n-3)}$ (fig. 6).






FIGURE 5. The path C_0 for Σ_7 . Double each polygon to obtain loops on Σ_7 , with punctures at vertices.


Any curve $\beta_{i,j}$ such that $i - j \equiv \pm 2 \pmod{n}$ bounds a twice punctured disk in Σ . There are once punctured annuli between the circles $\beta_{i,j}$ and $\beta_{i,j+1}$ or between $\beta_{i,j}$ and $\beta_{i+1,j}$. Create loops $B_{i,j}$ in $\Sigma_n \times S^1$, each corresponding to the circle $\beta_{i,j}$ in Σ_n , as in the Main Construction. Figures 7, 8, 9 show a picture of part of M_C for 5, 6, and 7 strand braids. The pictures are of a neighborhood of $\gamma \times I$, cut along a gap, and flattened out. Loops $B_{i,j}$ and $B_{i-1,j+1}$ have been isotoped to the same level.

A key observation is that if one does Dehn twists about the horizontal loops, *i.e.* by $1/n$ Dehn filling on these curves, then an incompressible surface which is obtained by tubing together pants

FIGURE 6. Half of the path C for Σ_5 FIGURE 7. M_C for Σ_5



in M may be isotoped across the loops without affecting its isotopy class in the Dehn filling. This follows since each pants is isotopic into a fiber surface, so the framing induced by the pants is the same as that of the fiber. For this reason, we may draw surfaces in M_C going through loops $B_{i,j}$ without drawing which side of the loop the surface lies on. Also, this makes it clear that two parallel pants may not be tubed together to get an incompressible surface.

We have a 2-complex in the trivial braid which consists of the loops and punctured disks and annuli (this is shown for $n = 5, 6, 7$ in figures 7, 8, 9). The intersection with $\gamma \times I$ is shown in figure

FIGURE 8. Part of M_C for Σ_6

10(a) for Σ_7 . Take the subcomplex consisting only of punctured annuli (fig. 10(b)). Then one can see that any surface carried by this complex is a punctured torus, isotopic to the punctured torus T which consists of the annuli bounded by the sequence $B_{1,3}, B_{1,4}, B_{2,4}, B_{2,5}, B_{3,5}, \dots, B_{n,3}, B_{1,3}$ (fig. 10(d)). If $n = 5$, then this is the only possibility for a surface. To see that every surface is isotopic to T if $n > 5$, note that the punctured annulus spanning $B_{i,j}, B_{i,j+1}, B_{i+1,j+1}$ is isotopic to the punctured annulus spanning $B_{i,j}, B_{i+1,j}, B_{i+1,j+1}$ (this follows from the symmetry of the A -regions: tubing two once punctured annuli gives a 4-punctured sphere, which is isotopic to the 4-punctured sphere on the other side of the A -region, as in figure 2). We may thus assume that the surface does not go through the loops $B_{i,j}$ where $i - j \pmod n = 2$ (fig. 10(b)). By induction, we may assume that for $i - j \pmod n \leq k$, the torus does not go through $B_{i,j}$, where $k < n - 3$ (fig. 10(c-d)).


Thus, we may assume that the surface goes through the sequence of loops $B_{i,j}$, where $i - j \pmod n = n - 3, n - 2$, which gives the punctured torus T . But T is pairwise compressible, which can be seen by considering the 2-punctured annulus spanning $\{B_{1,3}, B_{1,4}, B_{2,4}\}$ (fig. 11 gives a picture of the annular compression). This shows that there is no pairwise incompressible surface in T_ψ which is isotopic into M_C . If one performs Dehn twists about the curves $B_{i,j}$ avoiding the finite set of Lagrangian subspaces coming from Hatcher's theorem, one gets a small pure braid. In fact,

FIGURE 9. Part of M_C for Σ_7 FIGURE 10. The intersection of the complex with $\gamma \times I$ for Σ_7 .

one may get a link in S^3 by considering $n - 1$ strands as lying in a solid torus, and embedding this solid torus standardly in S^3 . $\square$

4. CONCLUSION

Given some natural complexity on 3-manifolds, such as the minimal number of tetrahedra in a triangulation, or the minimal number of intersections in a Heegaard diagram, it would be interesting to understand how the density of small 3-manifolds of complexity n behaves as $n \rightarrow \infty$, although this is probably a very difficult problem. A related question would be to consider all

FIGURE 11. An annular compression of T .

2-complexes coming from closed paths in $\mathcal{P}(\Sigma)^{(1)}$, and to consider the fraction of these for which the construction given in theorem 3.1 produces manifolds by Dehn twisting along the curves in the pants decompositions which have no incompressible surface which is isotopic off of the loops, for a given length of paths in the pants complex. We conjecture that this fraction goes to 1 as the length of the paths goes to ∞ . Indeed one can generalize the argument of theorem 3.1 to show that if a path goes through a subsequence of pants decompositions as defined in the proof of the theorem, then any closed incompressible surface in the complement of the pants curves will compress under Dehn twisting.

It should be possible to generalize the construction in this paper to prove the existence of knots in S^3 with arbitrarily large Heegaard genus. It would be interesting to understand this construction for surfaces with genus. One might be able to prove the existence of fibred manifolds with fiber of arbitrarily high genus, such that the fiber is the only closed connected incompressible surface.

REFERENCES

- [1] J. F. BROCK, *Weil-Petersson translation distance and volumes of mapping tori*, arXiv:math.GT/0109050.
- [2] D. COOPER AND M. SCHARLEMANN, *The structure of a solvmanifold's Heegaard splittings*, in Proceedings of 6th Gökova Geometry-Topology Conference, vol. 23, 1999, pp. 1–18.
- [3] M. CULLER, W. JACO, AND H. RUBINSTEIN, *Incompressible surfaces in once-punctured torus bundles*, Proc. London Math. Soc. (3), 45 (1982), pp. 385–419.
- [4] N. DUNFIELD, *Which small volume hyperbolic 3-manifolds are haken?* talk given at University of Warwick, slides available at http://www.math.harvard.edu/~nathand/preprints/slides/haken_slides.ps, July 1999.
- [5] W. FLOYD AND A. HATCHER, *Incompressible surfaces in punctured-torus bundles*, Topology Appl., 13 (1982), pp. 263–282.
- [6] W. HAKEN, *Some results on surfaces in 3-manifolds*, in Studies in Modern Topology, Math. Assoc. Amer. (distributed by Prentice-Hall, Englewood Cliffs, N.J.), 1968, pp. 39–98.
- [7] J. HASS AND W. MENASCO, *Topologically rigid non-Haken 3-manifolds*, J. Austral. Math. Soc. Ser. A, 55 (1993), pp. 60–71.
- [8] A. HATCHER, P. LOCHAK, AND L. SCHNEPS, *On the Teichmüller tower of mapping class groups*, J. Reine Angew. Math., 521 (2000), pp. 1–24.
- [9] A. HATCHER AND W. THURSTON, *Incompressible surfaces in 2-bridge knot complements*, Invent. Math., 79 (1985), pp. 225–246.
- [10] A. E. HATCHER, *On the boundary curves of incompressible surfaces*, Pacific J. Math., 99 (1982), pp. 373–377.

- [11] W. H. JACO AND P. B. SHALEN, *Seifert fibered spaces in 3-manifolds*, Mem. Amer. Math. Soc., 21 (1979), pp. viii+192.
- [12] L. M. LOPEZ, *Alternating knots and non-Haken 3-manifolds*, Topology Appl., 48 (1992), pp. 117–146.
- [13] ———, *Small knots in Seifert fibered 3-manifolds*, Math. Z., 212 (1993), pp. 123–139.
- [14] D. MARGALIT, *The automorphism group of the pants complex*, arXiv:math.GT/0201319.
- [15] J. W. MORGAN, *On Thurston’s uniformization theorem for three-dimensional manifolds*, in The Smith conjecture (New York, 1979), Academic Press, Orlando, FL, 1984, pp. 37–125.
- [16] Y. MORIAH AND H. RUBINSTEIN, *Heegaard structures of negatively curved 3-manifolds*, Comm. Anal. Geom., 5 (1997), pp. 375–412.
- [17] U. OERTEL, *Closed incompressible surfaces in complements of star links*, Pacific J. Math., 111 (1984), pp. 209–230.
- [18] A. THOMPSON, *Thin position and the recognition problem for S^3* , Math. Res. Lett., 1 (1994), pp. 613–630.
- [19] W. P. THURSTON, *The geometry and topology of 3-manifolds*. Lecture notes from Princeton University, 1978–80.
- [20] W. P. THURSTON, *Three-dimensional manifolds, Kleinian groups and hyperbolic geometry*, Bull. Amer. Math. Soc. (N.S.), 6 (1982), pp. 357–381.
- [21] F. WALDHAUSEN, *Recent results on sufficiently large 3-manifolds*, in Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976), Part 2, Amer. Math. Soc., Providence, R.I., 1978, pp. 21–38.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT CHICAGO, 322 SEO M/C 249, 851 S. MORGAN STREET,
CHICAGO, IL 60607-7045

E-mail address: agol@math.uic.edu